

Constraining the (γ, π) amplitude for E2 $N \rightarrow \Delta$

The ratio of the E2/M1 $N \rightarrow \Delta$ transitions provides an important discrimination between nucleon structure models. The quantity often compared to model calculations is the ratio of the corresponding photo-pion multipoles, $R_{EM}^{3/2} = \Im m(E_{1+}^{3/2})/\Im m(M_{1+}^{3/2})$, at the energy where the πN P_{33} phase goes through 90° . Recently, multipole analyses of two new data sets have reported new values for this quantity, $-2.5(\%) \pm 0.2(stat) \pm 0.2(sys)$ from Mainz [1], and $-2.9(\%) \pm 0.3(stat + sys) \pm 0.2(model)$ from LEGS [2]. Here we investigate the puzzling fact that while these multipole ratios appear to agree, the data from which they were extracted do not.

At least 8 observables are necessary to specify the photo-pion amplitude [3]. In the absence of such complete information, constraints from many observables are needed to avoid the potential ambiguity discussed by Donnachie [4] of higher partial wave strength appearing in lower partial waves, and vice versa. The Mainz analysis of [1] assumed complete dominance of S and P waves in a fit to only 2 observables. (In fact, the presence of an s -channel resonance guarantees non-Born u -channel contributions to higher partial waves that cannot *a priori* be ignored.) To examine the effect of these assumptions, we have repeated the analysis of the Mainz data, successively adding constraints from other observables and increasing the number of fitted partial waves. To avoid confusion, we have not included any LEGS data.

In [1], a fit of the Mainz 270-420 MeV $\sigma(\pi^\circ)$ cross section and $\Sigma(\pi^\circ)$ beam asymmetry data to powers of $\cos(\theta)$ was used to extract a quantity which, if only S and P waves contribute, should be the ratio of π° multipoles, $R_{\pi^\circ} = \Re[E_{1+}^{\pi^\circ} \cdot (M_{1+}^{\pi^\circ} - M_{1-}^{\pi^\circ})^*]/\{|E_{0+}^{\pi^\circ}|^2 + |3E_{1+}^{\pi^\circ} - (M_{1+}^{\pi^\circ} - M_{1-}^{\pi^\circ})|^2\}$. The authors argued that $R_{\pi^\circ}(340)$ is essentially $R_{EM}^{3/2}$. We have performed an equivalent energy-independent analysis of their data, fitting S and P-wave π° multipoles, with phases fixed to the elastic $\pi^\circ p$ values of the VPI[SM95] πN scattering solution [5]. The result, $R_{\pi^\circ} = -2.31(\%) \pm 0.54(stat + sys)$, is in agreement with the value of [1].

We now add constraints from other data. Following [2], we use a K-matrix-like parameterization in an energy-dependent analysis of the (γ, π) multipoles with isospin τ and orbital angular momentum ℓ , $A_{\ell\pm}^\tau(E_\gamma) = \{A_B^\tau(E_\gamma) + X(\varepsilon_\pi)\}(1 + iT_{\pi N}^\ell) + \beta \cdot T_{\pi N}^\ell$. Here, E_γ is the beam energy, A_B^τ is the pseudo-vector Born multipole (up to $\ell = 19$), and X is a polynomial in the π^+ kinetic energy, ε_π . VPI[SM95] values are used for the πN scattering T-matrix, $T_{\pi N}^\ell$ [5].

The isospin decomposition requires π^+ data. For this we use $\sigma(\pi^+)$ from [6] and $\Sigma(\pi^+)$ from [7]. Varying S and P waves gives $-(3.27 \pm 0.35)\%$. In row 1 of the table we show the effect of extending the fit to higher partial waves. The result, constrained by only 2 observables, suffers from Donnachie's ambiguity [4] and is not stable.

In row 2 we add constraints from other polarization ratios, the Bonn target (T) asymmetries [8], and Khar'kov target and recoil (P) asymmetries [7,9].

The Mainz π^0 cross sections of [1] are in very good agreement with the Bonn results from [10]. However, the older Bonn data cover a larger angular range. Including these in the fit gives the third row of the table. The result is finally independent of the number of partial waves, but is significantly lower than R_{π^0} from [1].

In conclusion, the procedure in [1] of fitting only S and P-wave multipoles to only 2 observables essentially corresponds to truncating an expansion before it has converged. Other authors, refs. [11,12], have disputed the accuracy of approximating $R_{\text{EM}}^{3/2}$ with R_{π^0} . While we agree that this approximation is quite sensitive to the multipole solution, such inaccuracies are in fact completely dwarfed by ambiguities in multipoles fitted with the Mainz procedure. This work was supported by U.S. Dept. of Energy Contract No. AC02-76CH00016.

A.M. Sandorfi¹, J. Tonnison^{2,1} and S. Hoblit¹

¹Physics Dept., Brookhaven National Lab, Upton, NY 11973

²Physics Dept., Virginia Polytechnic Inst. and State Univ., Blacksburg, Va 24061

Received

PACS numbers: 11.80.Et, 14.20.Gk, 13.60.Le

REFERENCES

- [1] R. Beck *et al.*, Phys. Rev. Lett. **78**, 606, 1997; H.-P. Krahn, thesis, U. Mainz (1996).
- [2] LEGS Collaboration, Phys. Rev. Lett. (submitted); BNL-64382, from <http://www.legs.bnl.gov/>.
- [3] W. Chiang and F. Tabakin, Phys. Rev. **C55**, 2054 (1997).
- [4] A. Donnachie, Rep. Prog. Phys. **36**, 695 (1973).
- [5] R. Arndt and R.L. Workman, SAID code, telnet vtinte.phys.vt.edu [*physics:quantum*].
- [6] G. Fischer *et al.*, Z. Physik **253**, 38 (1972).
- [7] V.A. Get'man *et al.*, Nucl. Phys. **B188**, 397 (1981).
- [8] H. Dutz *et al.*, Nucl. Phys. **A601**, 319, (1996); Gisela Anton, priv. comm.
- [9] A. Belyaev *et al.*, Nucl. Phys. **B213**, 201 (1983).
- [10] H. Genzel *et al.*, Z. Physik **268**, 43 (1974).
- [11] R. Davidson and N. Mukhopadhyay, Nucl-th. 9705035.
- [12] R.L. Workman, Nucl-th. 9702052.

TABLES

TABLE I. Variation of $R_{\text{EM}}^{3/2}$ with increasing number of fitted partial waves, in successive columns. Constraints from different observables between 270-420 MeV are increased in successive rows: row 1, refs. [1,6,7]; row 2, refs. [1,6–9]; row 3: refs. [1,6–10].

Data used in	$R_{\text{EM}}^{3/2}(\%)$	$R_{\text{EM}}^{3/2}(\%)$	$R_{\text{EM}}^{3/2}(\%)$
analysis	$\ell_\pi = S - P$	$\ell_\pi = S - D$	$\ell_\pi = S - F$
σ, Σ	$-(3.27 \pm 0.35)$	$-(4.75 \pm 0.56)$	$-(5.66 \pm 0.63)$
σ, Σ, T, P	$-(1.40 \pm 0.24)$	$-(2.93 \pm 0.30)$	$-(3.39 \pm 0.38)$
$\sigma, \Sigma, T, P, \sigma$	$-(0.58 \pm 0.08)$	$-(0.71 \pm 0.10)$	$-(0.58 \pm 0.12)$